

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

SDSU Extension Fact Sheets

SDSU Extension

1960

An All-Pullet Flock for More Profit

Cooperative Extension

South Dakota State University

Follow this and additional works at: https://openprairie.sdstate.edu/extension_fact

Recommended Citation

Extension, Cooperative, "An All-Pullet Flock for More Profit" (1960). *SDSU Extension Fact Sheets*. 194.
https://openprairie.sdstate.edu/extension_fact/194

This Fact Sheet is brought to you for free and open access by the SDSU Extension at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in SDSU Extension Fact Sheets by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

Historical, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



For current policies and practices, contact SDSU Extension

Website: extension.sdstate.edu

Phone: 605-688-4792

Email: sdsu.extension@sdstate.edu

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.

Thistles

CANADA THISTLE¹ and perennial sowthistle² are deep-rooted perennials that spread by seeds and underground parts. They emerge later in the spring and are less difficult to kill than leafy spurge, Russian knapweed or hoary cress. They emerge about the same time as bindweed.

They are cross-pollinated so flowers must be open before seed can be produced. Seeds are developed early and are ready to germinate 8 to 10 days after the flowers have opened. Each seed is attached to a tiny parachute that can be carried miles by air currents.

Single plants spread by means of seeds and roots to develop patches like other perennial weeds. Seeds scattered by the wind develop into sprinklings of plants throughout the countryside, which creates a different problem than that associated with other noxious weeds. These light infestations sometimes are not recognized as a crop hazard. They do reduce yields, however. In one test 2 Canada thistle plants per square yard reduced wheat yield 18% while 19 plants per square yard reduced yields 36%. A patch of sowthistle caused a 69% reduction of oats yields.

To control or eliminate these thistles, use intensive cultivation, soil sterilant chemicals, certain competitive crops, selective herbicides, or several combinations of cultivation, crops and chemicals.

INTENSIVE CULTIVATION

Intensive cultivation from spring until freeze-up will kill a high percentage of the thistles. Cultivation from immediately after harvest one year until freeze-up the next is more effective.

A duckfoot cultivator or one-way disk are equally satisfactory implements. However, it is frequently desirable to perform the first operation with a plow if there is considerable plant residue. Equip the duckfoot cultivator with wide sweeps (12 to 24 inches) that overlap 3 or 4 inches. Keep them sharp and see that they are flat when in the soil and operating at a depth of 4 to 5 inches. The same is true for the one-way disk—keep the disks sharp and operate at a depth of 4 to 5 inches. It is essential to cut each thistle root at each cultivation.

It takes 10 to 15 days for the plants to emerge after the roots have been cut. Another 10 to 15 days elapse before there are enough leaves to produce more food than is needed for growth. Therefore, little food is stored in the roots and root reserves are being used for plant growth for a period of 20 to 30 days. Each cultivation has a similar effect. Cultivate every 3 weeks during good growing conditions and every 4 weeks during periods of dry, hot weather when the plants are growing less rapidly. This generally means that cultivations should be done at 3-week intervals during June and July and at 4-week intervals during August, September and October.

A disk harrow is also a satisfactory implement. Experiments have shown that double disking with a tandem disk was satisfactory when used every 2 weeks (at least 16 operations) on plowing.

Combining intensive cultivation for part of the season with the production of a crop and a chemical application is generally more practical than an entire season of cultivation. Income from the crop is obtained and the hazards of erosion, resulting from a full season of cultivation, are greatly reduced.

SPRAYING

There are several chemicals that can be used for controlling thistles. For controlling sowthistle, 2,4-D is generally more effective than 2,4,5-T or MCPA; however, MCPA is often equal to and sometimes superior to 2,4-D on Canada thistle. An ester formulation of 2,4-D is sometimes superior to an amine formulation for controlling sowthistle, but there is

Thistles

Control and Elimination of

Cooperative Extension Service



F.S. 20

seldom much difference between the two on Canada thistle.

Use one-third to one-half pound acid equivalent per acre of 2,4-D or MCPA to kill seedlings, prevent seed production on mature plants, kill some old plants and weaken others so that later treatments kill them. Treat when the weeds are in the rosette stage of growth (either spring or fall). Two treatments in 1 year (spring and fall rosettes) will kill up to 85% of the thistles. As a general rule, however, thistles are not killed by two treatments the first year are more resistant to the chemical and cannot be killed by spraying. Some other treatment, such as cultivation, is needed to eliminate them.

Amritol (sometimes called ATA, amizol, or amino triazole) generally kills over 90% of Canada thistles. Apply at the rate of 4 to 6 pounds acid equivalent in 10 to 30 gallons of water per acre when the thistles are starting to bud. Similar treatments made to thistles 6 to 8 inches tall generally give good top kill but very little root kill. Excellent results are sometimes obtained by treating regrowth of thistles that have been mowed or plowed. Treatments should be made with 4 to 6 pounds acid equivalent per acre after regrowth is 8 inches tall and before buds start.

Amritol is not a selective spray and will injure most crops. It acts slowly and thistles may not be completely dead for 6 weeks. Thistles droop and turn yellow in about 10 days; however, they sometimes appear to recover by the end of 3 or

Canada Thistle



Perennial Sow Thistle



By Lyle A. Derscheld, Experiment Station Agronomist, and
Keith E. Wallace, Extension Weed Specialist
(Experiment Station Project 32-R)

¹*Cirsium arvense* L.

²*Sonchus oleraceus* L. and *S. uliginosus* Bieb.

4 weeks but then die. Regrowth will generally be yellow. Amritol is somewhat expensive but is frequently useful for eliminating patches. It is generally less effective on sow-thistle than on Canada thistle.

One to 2 pounds acid equivalent per acre of 2,4-DB frequently prevent seed production of Canada thistle and weaken the plants. Because of its cost, its use will probably be limited to controlling broadleaved weeds in new seedings or alfalfa. However, it may be useful in checking Canada thistle until the legume can become established.

CULTIVATION, CROPS, CHEMICALS

Small grains. A high percentage of thistles can be eliminated in a year when small grain is being produced. Since barley and wheat are generally more tolerant than oats to 2,4-D, they are usually better adapted for this use. Spray the grain with one-half to three-fourths pound acid equivalent of 2,4-D amine or MCPA amine per acre about June 1 when the thistles are not over 6 inches tall and the grain is in the 5- to 6-leaf stage of growth. Such a treatment will prevent the production of seed and will weaken the plants. Follow-up treatments after harvest will deliver the knockout punch.

Experimental results indicate that the percentage of thistles killed will depend on the type of follow-up treatment used.

(1) Three cultivations after harvest (one plowing and two cultivations with duckfoot cultivator or one-way disk) killed 85% of the weeds the first year. A second year of spraying in the grain and three cultivations after harvest gave 100% elimination.

(2) Plowing after harvest and spraying thistles with three-fourths pound 2,4-D or MCPA acid equivalent per acre after they emerged on the plowing killed 85% of the weeds in 1 year. A second year of spraying in the grain, plowing after harvest, and spraying after thistles emerged on the plowing gave 98% elimination.

(3) Spraying in the stubble after harvest gave 50% kill. Three years of spraying in the grain and again after harvest gave over 90% elimination.

(4) Late fall plowing (late October) gave 70% kill in 3 years when used as the only fall treatment after spraying in the grain each year. However, it did not improve results when any of the fall treatments listed were used.

One application of 2,4-D or MCPA in corn kills very few thistles. However, two applications of one-half to three-fourths pound per acre (once before the second cultivation and once after tasseling with drop nozzles) can be expected to kill about 40% of the weeds. Sprayed weeds should not be cultivated for about 1 week after spraying.

One season of intensive cultivation followed by a crop of winter wheat or rye will kill a high percentage of thistles in 1 year and keep the soil from eroding over winter. Likewise, complete elimination can usually be achieved in 2 or 3 years by growing a crop of winter wheat or rye and cultivating every 3 or 4 weeks between harvest and seeding.

Summer Crops. Two or three cultivations, followed by a late June seeding of soybeans, sorghum, or sudan grass with a grain drill, generally eliminate a high percentage of thistles in 1 year. The crop should be harvested for forage and the area cultivated twice after harvest to get the best results. At least two crops and sometimes three are generally required to give complete kill of all thistles.

Forage Crops. A good crop of alfalfa or alfalfa-bromegrass mixture, which is cut regularly for hay, almost eliminates thistles in 3 years. The use of 2,4-DB may be useful for controlling annual broad-leaved weeds and Canada thistle the first year when establishing the stand. Likewise dalapon at the rate of 2 pounds acid equivalent per acre may be used to control annual grassy weeds in alfalfa the first year. If either chemical is used, however, hay produced the first year should not be fed to livestock.

A good crop of bromegrass which is cut regularly for hay will generally give over 90% elimination in 3 years. Spraying

with 2,4-D twice a year (early June and mid-August) in a good stand of bromegrass, crested wheatgrass, or other strongly competitive perennial grass will generally kill over 90% of the thistles in 3 years. Use three-fourths pound acid equivalent of the 2,4-D, except use only one-half pound the spring that the grass is seeded.

Elimination of thistles can be achieved in less time by intensively cultivating the infested area five times prior to the seeding of a forage crop in mid-August. About 90% elimination can be obtained the first year and 95% the second year. However, the use of 2,4-D in grasses the second year will generally give 100% elimination by the end of the second year.

Fallow. In cases where a 3- or 4-year rotation includes a year of fallow, there are several methods that can be used during the fallow year to substantially reduce the stand of Canada thistle.

One system uses one spring cultivation with duckfoot cultivator or one-way disk, a spraying when thistles are starting to produce buds, and continued cultivation 6 weeks after spraying. Either one pound acid equivalent of 2,4-D, 3 pounds of amritol or 4 pounds of 2,4-DB per acre may be used as the spray.

A second method includes a spring cultivation, an application of $\frac{1}{2}$ to $\frac{3}{4}$ pound of 2,4-D per acre when the thistles are starting to bud, a second application of $\frac{1}{2}$ to $\frac{3}{4}$ pound of 2,4-D two weeks later, and a late fall cultivation just before freezeup in October.

SOIL STERILANTS

Numerous soil sterilants can be used to eliminate patches of thistles with one treatment. Most of the chemicals give best results when applied between September 1 and December 1; however, good results are often obtained from summer applications. Apply the chemical to a band 6 or 8 feet wide around the outside of the patch to kill roots that extend beyond the patch.

The following chemicals are generally effective in giving at least 95% elimination when applied at the rates designated for each square rod. The lower rates are satisfactory for fall treatments, but the higher rates are sometimes needed for summer applications.

Brand Name	Rate per Square Rod	Cost*
Ammate "X"	4 to 6 pounds	\$1.40
Atlacide	6 to 8 pounds	1.08
Chlorax	10 to 12 pounds	3.20
Chlorox	6 to 8 pounds	1.90
Concentrated Borascu	12 to 15 pounds	0.85
DB Granular	5 to 6 pounds	0.95
Novon Concentrate	$\frac{1}{2}$ to 1 quart	0.76
Polybor-chlorate	10 to 12 pounds	1.75
Sodium chlorate	5 pounds	0.90

*Approximate retail cost of chemical for 1 square rod at lower rate for areas 10 to 15 square rods in size. Cost decreases for large areas and varies somewhat with local situations.

CONTROL SEED PRODUCTION

Prevention of seed production is more important for thistles than for most noxious weeds. It can be done by mowing or spraying at the proper time. Since a high percentage of seeds are ready to germinate after flowers have been open 8 to 10 days, mowing must be done during the week after they bloom. Likewise, spraying with one-third to one-half pound of 2,4-D must be done before blooming.

You need not be concerned about thistle buds in seed grain. They must open before pollination takes place and they must be open over a week before seeds are mature enough to germinate. Although there is no possibility of the buds containing viable seeds, there is still a possibility that older heads were threshed. There may be individual seeds in the grain.